

Quasiconvex stochastic processes and a separation theorem

DAWID KOTRYS AND KAZIMIERZ NIKODEM

Dedicated to Professor János Aczél on his 90th birthday

Abstract. Quasiconvex stochastic processes are introduced. A characterization of pairs of stochastic processes that can be separated by a quasiconvex stochastic process and a stability theorem for quasiconvex processes are given.

Mathematics Subject Classification (2010). Primary: 26A51 · Secondary: 60G99.

Keywords. Quasiconvex stochastic process · Separation by quasiconvex stochastic processes · Stability of quasiconvex stochastic processes.

1. Introduction

In [1], Baron et al. proved that two real functions f and g defined on a real interval I can be separated by a convex function if and only if they fulfil the following inequality

$$f(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y),$$

for all $x, y \in I$ and $\lambda \in [0, 1]$.

In 1994 Smolarz [11] obtained an analogous result for quasiconvex functions. Namely, he proved that two functions $f, g : I \rightarrow \mathbb{R}$ can be separated by a quasiconvex function if and only if

$$f(\lambda x + (1 - \lambda)y) \leq \max\{g(x), g(y)\},$$

for all $x, y \in I$ and $\lambda \in [0, 1]$ (see also [3]).

In this paper we introduce the notion of quasiconvex stochastic processes and present some properties of them. In particular we show that a stochastic process is convex if and only if it is Jensen-convex and quasiconvex. Our main result extends the Smolarz separation theorem to quasiconvex stochastic

processes. As a consequence we obtain a Hyers-Ulam-type stability result for quasiconvex stochastic processes.

2. Preliminaries

Let (Ω, \mathcal{A}, P) be an arbitrary probability space and $I \subset \mathbb{R}$ be an interval. A function $X : \Omega \rightarrow \mathbb{R}$ is called a *random variable*, if it is \mathcal{A} -measurable. A function $X : I \times \Omega \rightarrow \mathbb{R}$ is called a *stochastic process*, if for every $t \in I$ the function $X(t, \cdot)$ is a random variable.

Recall that a stochastic process $X : I \times \Omega \rightarrow \mathbb{R}$ is said to be *convex*, if

$$X(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \lambda X(t_1, \cdot) + (1 - \lambda)X(t_2, \cdot) \quad (\text{a.e.}),$$

for all $t_1, t_2 \in I$ and $\lambda \in [0, 1]$.

A stochastic process $X : I \times \Omega \rightarrow \mathbb{R}$ is called *Jensen-convex*, if

$$X\left(\frac{t_1 + t_2}{2}, \cdot\right) \leq \frac{X(t_1, \cdot) + X(t_2, \cdot)}{2} \quad (\text{a.e.}),$$

for all $t_1, t_2 \in I$.

Convex and Jensen-convex stochastic processes were investigated by many authors and various properties and applications of them can be found in the literature (see, for instance, [5, 6, 9, 10] and the references therein).

We say that a stochastic process $X : I \times \Omega \rightarrow \mathbb{R}$ is *quasiconvex*, if

$$X(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \max\{X(t_1, \cdot), X(t_2, \cdot)\} \quad (\text{a.e.}), \quad (1)$$

for all $t_1, t_2 \in I$ and $\lambda \in [0, 1]$.

We start our investigation with two simple observations.

Observation 1. *The following conditions are equivalent:*

- (i) $X : I \times \Omega \rightarrow \mathbb{R}$ is a quasiconvex stochastic process.
- (ii) For every random variable $A : \Omega \rightarrow \mathbb{R}$ the level set

$$L_A = \{t \in I : X(t, \cdot) \leq A(\cdot)\} \quad (\text{a.e.})$$

is convex.

Proof. Suppose first that X is a quasiconvex stochastic process and fix a random variable $A : \Omega \rightarrow \mathbb{R}$. Let $t_1, t_2 \in L_A$ and $\lambda \in [0, 1]$. By (1) and the definition of level sets we have

$$\begin{aligned} X(\lambda t_1 + (1 - \lambda)t_2, \cdot) &\leq \max\{X(t_1, \cdot), X(t_2, \cdot)\} \\ &\leq \max\{A(\cdot), A(\cdot)\} = A(\cdot) \quad (\text{a.e.}). \end{aligned}$$

Thus $\lambda t_1 + (1 - \lambda)t_2 \in L_A$, which proves that L_A is convex.

Assume now that the sets L_A are convex for all random variables A . Fix $t_1, t_2 \in I$ and $\lambda \in [0, 1]$. Define $A(\cdot) = \max\{X(t_1, \cdot), X(t_2, \cdot)\}$. Then, of course,

$t_1, t_2 \in L_A$, and, by the convexity of L_A , we have $\lambda t_1 + (1 - \lambda)t_2 \in L_A$. It means that the inequality

$$X(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq A(\cdot) = \max\{X(t_1, \cdot), X(t_2, \cdot)\} \quad (\text{a.e.})$$

holds and X is quasiconvex. \square

Observation 2. *If a stochastic process $X : I \times \Omega \rightarrow \mathbb{R}$ is convex, then it is quasiconvex.*

Proof. By the convexity of X , for all $t_1, t_2 \in I$ and $\lambda \in [0, 1]$, we have

$$\begin{aligned} X(\lambda t_1 + (1 - \lambda)t_2, \cdot) &\leq \lambda X(t_1, \cdot) + (1 - \lambda)X(t_2, \cdot) \\ &\leq \max\{X(t_1, \cdot), X(t_2, \cdot)\} \quad (\text{a.e.}), \end{aligned}$$

which shows that the process X is quasiconvex. \square

Clearly, quasiconvex (as well as Jensen-convex) stochastic processes need not be convex. However, if a stochastic process is both quasiconvex and Jensen-convex, then it is convex.

Proposition 3. *Let I be an open interval. A stochastic process $X : I \times \Omega \rightarrow \mathbb{R}$ is convex if and only if it is quasiconvex and Jensen-convex.*

Proof. The “only if” part is clear. To prove the “if” part fix $t_1, t_2 \in I$, $t_1 < t_2$. By the quasiconvexity of X , for every $t \in [t_1, t_2]$, we have

$$X(t, \cdot) \leq \max\{X(t_1, \cdot), X(t_2, \cdot)\} \quad (\text{a.e.}).$$

This implies that the process X is P -upper bounded on $[t_1, t_2]$, that is

$$\lim_{n \rightarrow \infty} \sup_{t \in [t_1, t_2]} \{P(\{\omega \in \Omega : |X(t, \omega)| \geq n\})\} = 0.$$

Since X is also Jensen-convex, it follows, by the Bernstein-Doetsch-type theorem, that X is continuous in probability and, consequently, convex (see [6] Theorems 4, 5). \square

3. Main result

At the beginning of this section we would like to recall the definition and basic properties of the essential infimum of a collection of functions. Let $(\Omega, \mathcal{F}, \mu)$ be a measure space and \mathcal{S} be a collection of measurable functions $f : \Omega \rightarrow \mathbb{R}$. On \mathbb{R} the Borel σ -algebra is used. If \mathcal{S} is a countable set, then we may define the pointwise infimum of the functions from \mathcal{S} , which will itself be measurable. If \mathcal{S} is uncountable, then the pointwise infimum need not be measurable. In this case, the essential infimum can be used. The *essential infimum* of \mathcal{S} , written as $\text{ess inf } \mathcal{S}$, if it exists, is a measurable function $f : \Omega \rightarrow \mathbb{R}$ satisfying the two following axioms:

- $f \leq g$ almost everywhere, for any $g \in \mathcal{S}$,

- if $h : \Omega \rightarrow \mathbb{R}$ is measurable and $h \leq g$ almost everywhere for every $g \in \mathcal{S}$, then $h \leq f$ almost everywhere.

Note that if f is the essential infimum and $g : \Omega \rightarrow \mathbb{R}$ is equal to f almost everywhere, then g is also an essential infimum. Conversely, if f and g are both essential infima, then, from the above definition $f \leq g$ and $g \leq f$, so $f = g$ almost everywhere. It can be shown that for a σ -finite measure μ , the essential infimum of \mathcal{S} does exist. Furthermore, there exists a sequence $(f_n)_{n \in \mathbb{N}}$ in \mathcal{S} such that

$$\text{ess inf } \mathcal{S} = \inf \{f_n : n \in \mathbb{N}\}.$$

For more details we refer the reader to [2].

The following properties of essential infimum will be useful in the sequel.

Lemma 4. *Let $(\Omega, \mathcal{F}, \mu)$ be a σ -finite measure space, \mathcal{S} be a nonempty collection of measurable real functions defined on Ω , and let $g : \Omega \rightarrow \mathbb{R}$ be a measurable function. If $\text{ess inf } \mathcal{S} < g$ almost everywhere, then there exist sets $\Omega_n \in \mathcal{F}$ and functions $f_n \in \mathcal{S}$ for $n \in \mathbb{N}$, such that $\mu(\Omega \setminus \bigcup_{n \in \mathbb{N}} \Omega_n) = 0$ and $f_n < g$ on Ω_n , $n \in \mathbb{N}$.*

Proof. By the fact mentioned above there exists a sequence $(f_n)_{n \in \mathbb{N}}$ of elements of \mathcal{S} such that

$$\text{ess inf } \mathcal{S} = \inf \{f_n : n \in \mathbb{N}\}.$$

Assume that $\inf \{f_n : n \in \mathbb{N}\} < g$ on $\bar{\Omega} = \Omega \setminus \Omega_0$, where $\mu(\Omega_0) = 0$. By the definition of the infimum, for every $\omega \in \bar{\Omega}$ there exists $n \in \mathbb{N}$ such that $f_n(\omega) < g(\omega)$.

Define the sets

$$\Omega_n = \{\omega \in \bar{\Omega} : f_n(\omega) < g(\omega)\}.$$

Then $\Omega_n \in \mathcal{F}$, $\bigcup_{n \in \mathbb{N}} \Omega_n = \bar{\Omega}$, and $f_n < g$ on Ω_n for $n \in \mathbb{N}$, which was to be proved. \square

Lemma 5. *Let $f : \Omega \rightarrow \mathbb{R}$ be a measurable function and \mathcal{S} be a family of all measurable functions $g : \Omega \rightarrow \mathbb{R}$ such that $f < g$ almost everywhere. Then $\text{ess inf } \mathcal{S} = f$.*

Proof. Clearly $\text{ess inf } \mathcal{S} \geq f$ almost everywhere. To conclude the proof it is enough to observe that $f + \frac{1}{n} \in \mathcal{S}$ for all $n \in \mathbb{N}$. \square

Now we present our main theorem. It gives a condition under which two stochastic processes can be separated by a quasiconvex stochastic process.

Theorem 6. *Let $X, Y : I \times \Omega \rightarrow \mathbb{R}$ be stochastic processes. There exists a quasiconvex stochastic process $H : I \times \Omega \rightarrow \mathbb{R}$ such that*

$$X(t, \cdot) \leq H(t, \cdot) \leq Y(t, \cdot) \quad (\text{a.e.}), \quad t \in I,$$

if and only if

$$X(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \max\{Y(t_1, \cdot), Y(t_2, \cdot)\} \quad (\text{a.e.}) \quad (2)$$

for all $t_1, t_2 \in I$ and $\lambda \in [0, 1]$.

Proof. The sufficiency is obvious. To prove the necessity assume that X and Y fulfil (2). Given a random variable $A : \Omega \rightarrow \mathbb{R}$ consider the level set

$$L_A = \{t \in I : Y(t, \cdot) \leq A(\cdot) \quad (\text{a.e.})\}.$$

Let $C_A = \text{conv } L_A$ denote the convex hull of the set L_A . Define a stochastic process $H : I \times \Omega \rightarrow \mathbb{R}$ by

$$H(t, \cdot) = \text{ess inf}\{A : t \in C_A\}.$$

Fix $t_0 \in I$ and take a random variable $A : \Omega \rightarrow \mathbb{R}$ such that $t_0 \in C_A$. In view of the Caratheodory theorem (cf. [8]) we have $t_0 = \lambda t_1 + (1 - \lambda)t_2$, for some $t_1, t_2 \in L_A$ and $\lambda \in [0, 1]$. Hence $Y(t_1, \cdot) \leq A(\cdot) \quad (\text{a.e.})$ and $Y(t_2, \cdot) \leq A(\cdot) \quad (\text{a.e.})$ and, by inequality (2), we obtain

$$X(t_0, \cdot) = X(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \max\{Y(t_1, \cdot), Y(t_2, \cdot)\} \leq A(\cdot) \quad (\text{a.e.}).$$

Since the above inequality holds for any random variable A , such that $t_0 \in C_A$, we get

$$X(t_0, \cdot) \leq \text{ess inf}\{A : t_0 \in C_A\} = H(t_0, \cdot) \quad (\text{a.e.}).$$

Moreover, since for every fixed $t_0 \in I$ we have $t_0 \in L_{Y(t_0, \cdot)} \subset C_{Y(t_0, \cdot)}$, we get also

$$H(t_0, \cdot) = \text{ess inf}\{A : t_0 \in C_A\} \leq Y(t_0, \cdot) \quad (\text{a.e.}).$$

Now we will show that the stochastic process H is quasiconvex. Fix $t_1, t_2 \in I$ and $\lambda \in [0, 1]$. The following cases are possible:

- (i) $H(t_1, \cdot) \leq H(t_2, \cdot) \quad (\text{a.e.})$.
- (ii) $H(t_1, \cdot) > H(t_2, \cdot) \quad (\text{a.e.})$.
- (iii) The following sets Ω_1 and Ω_2 have positive measure

$$\begin{aligned} \Omega_1 &= \{\omega : H(t_1, \omega) \leq H(t_2, \omega)\}, \\ \Omega_2 &= \{\omega : H(t_1, \omega) > H(t_2, \omega)\}. \end{aligned}$$

Assume first that case (i) holds. We will show that there exist sets $\Omega_{n,k} \in \mathcal{A}$, $n, k \in \mathbb{N}$, such that $P(\bigcup_{n,k \in \mathbb{N}} \Omega_{n,k}) = 1$ and for all $n, k \in \mathbb{N}$, $H(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq H(t_2, \cdot)$ almost everywhere on $\Omega_{n,k}$.

Take an arbitrary random variable $B : \Omega \rightarrow \mathbb{R}$ satisfying

$$H(t_2, \cdot) = \text{ess inf}\{A : t_2 \in C_A\} < B(\cdot) \quad (\text{a.e.}) \quad (3)$$

By Lemma 4 there exist sets $\Omega_n \in \mathcal{A}$ and random variables $A_n \in \{A : t_2 \in C_A\}$ such that $P(\bigcup_{n \in \mathbb{N}} \Omega_n) = 1$ and for every $n \in \mathbb{N}$, $A_n < B$ on Ω_n . Then

$$\begin{aligned} t_2 \in C_{A_n} &= \text{conv}\{t \in I : Y(t, \cdot) \leq A_n(\cdot) \quad (\text{a.e.})\} \\ &\subset \text{conv}\{t \in I : Y(t, \cdot) < B(\cdot) \quad (\text{a.e.}) \quad \text{on} \quad \Omega_n\}. \end{aligned} \quad (4)$$

Now, we use the fact that $H(t_1, \cdot) \leq H(t_2, \cdot) < B(\cdot) \quad (\text{a.e.})$ and apply Lemma 4 separately for every set Ω_n , $n \in \mathbb{N}$. There exist sets $\Omega_{n,k} \subset \Omega_n$, $\Omega_{n,k} \in \mathcal{A}$ and random variables $A_{n,k} \in \{A : t_1 \in C_{A|_{\Omega_n}}\}$ such that $P(\bigcup_{k \in \mathbb{N}} \Omega_{n,k}) = P(\Omega_n)$ and $A_{n,k} < B$ on $\Omega_{n,k}$ for $n, k \in \mathbb{N}$. Then

$$\begin{aligned} t_1 \in C_{A_{n,k}|_{\Omega_n}} &= \text{conv}\{t \in I : Y(t, \cdot) \leq A_{n,k}(\cdot) \quad (\text{a.e.}) \quad \text{on} \quad \Omega_n\} \\ &\subset \text{conv}\{t \in I : Y(t, \cdot) < B(\cdot) \quad (\text{a.e.}) \quad \text{on} \quad \Omega_{n,k}\}. \end{aligned} \quad (5)$$

By (4) and (5), for every $n, k \in \mathbb{N}$ we obtain

$$t_1, t_2 \in \text{conv}\{t \in I : Y(t, \cdot) < B(\cdot) \quad (\text{a.e.}) \quad \text{on} \quad \Omega_{n,k}\}$$

and consequently

$$\lambda t_1 + (1 - \lambda)t_2 \in \text{conv}\{t \in I : Y(t, \cdot) < B(\cdot) \quad (\text{a.e.}) \quad \text{on} \quad \Omega_{n,k}\}.$$

Hence, by the definition of H , we get $H(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq B(\cdot) \quad (\text{a.e.})$ on $\Omega_{n,k}$. Since the family $(\Omega_{n,k})_{n,k \in \mathbb{N}}$ is countable and $P(\bigcup_{k \in \mathbb{N}} \Omega_{n,k}) = P(\Omega_n)$ for $n \in \mathbb{N}$ and $P(\bigcup_{n \in \mathbb{N}} \Omega_n) = 1$, we also have $H(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq B(\cdot) \quad (\text{a.e.})$ on Ω . Using the fact, that this inequality holds for every random variable B satisfying (3), by Lemma 5 we obtain

$$H(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq H(t_2, \cdot) = \max\{H(t_1, \cdot), H(t_2, \cdot)\} \quad (\text{a.e.}).$$

This finishes the the proof in case (i).

The proof in case (ii) is analogous, so we omit it.

Now assume that case (iii) holds. We consider two processes H_1 and H_2 being the restrictions of H to $I \times \Omega_1$ and $I \times \Omega_2$, respectively. Then for H_1 case (i) occurs and for H_2 case (ii) occurs. It means that

$$H_1(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \max\{H_1(t_1, \cdot), H_1(t_2, \cdot)\} \quad (\text{a.e.}) \quad \text{on} \quad \Omega_1$$

and

$$H_2(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \max\{H_2(t_1, \cdot), H_2(t_2, \cdot)\} \quad (\text{a.e.}) \quad \text{on} \quad \Omega_2.$$

Consequently

$$H(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \max\{H(t_1, \cdot), H(t_2, \cdot)\} \quad (\text{a.e.}) \quad \text{on} \quad \Omega$$

and the proof is complete. \square

4. Hyers–Ulam stability

As a direct consequence of Theorem 6 we obtain the following Hyers–Ulam-type stability result for quasiconvex stochastic processes. For the classical Hyers–Ulam theorem see [4]. The stability theorem for quasiconvex functions was obtained in [7] (cf. also [11]).

Theorem 7. *Let ε be a positive constant. If a stochastic process $X : I \times \Omega \rightarrow \mathbb{R}$ satisfies the inequality*

$$X(\lambda t_1 + (1 - \lambda)t_2, \cdot) \leq \max\{X(t_1, \cdot), X(t_2, \cdot)\} + \varepsilon \quad (\text{a.e.})$$

for all $t_1, t_2 \in I$ and $\lambda \in [0, 1]$, then there exists a quasiconvex stochastic process $H : I \times \Omega \rightarrow \mathbb{R}$ such that $|X(t, \cdot) - H(t, \cdot)| \leq \frac{\varepsilon}{2}$ (a.e.) for every $t \in I$.

Proof. To prove the above theorem it is enough to apply Theorem 6 to the stochastic processes X and $X + \varepsilon$. Hence, there exists a process $H_1 : I \times \Omega \rightarrow \mathbb{R}$, which is quasiconvex and satisfies $X(t, \cdot) \leq H_1(t, \cdot) \leq X(t, \cdot) + \varepsilon$ (a.e.). By taking $H(t, \cdot) = H_1(t, \cdot) - \frac{\varepsilon}{2}$ we get $|X(t, \cdot) - H(t, \cdot)| \leq \frac{\varepsilon}{2}$ (a.e.). This completes the proof. \square

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- [1] Baron, K., Matkowski, J., Nikodem, K.: A sandwich with convexity. *Math. Pannonica* **5/1**, 139–144 (1994)
- [2] Doob, J.L.: *Measure theory*. Graduate Texts in Mathematics. Springer, Berlin (1993)
- [3] Förg-Rob, W., Nikodem, K., Páles, Zs.: Separation by monotonic functions. *Math. Pannonica* **7/2**, 191–196 (1993)
- [4] Hyers, D.H., Ulam, S.M.: Approximately convex functions. *Proc. Am. Math. Soc.* **3**, 821–828 (1952)
- [5] Kotrys, D.: Hermite–Hadamard inequality for convex stochastic processes. *Aequat. Math.* **83**, 143–151 (2012)
- [6] Nikodem, K.: On convex stochastic processes. *Aequat. Math.* **20**, 184–197 (1980)
- [7] Nikodem, K.: Approximately quasiconvex functions. *C. R. Math. Rep. Acad. Sci. Canada* **10**, 291–294 (1988)
- [8] Roberts, A.W., Varberg, D.E.: *Convex functions*. Academic Press, New York (1973)
- [9] Shaked, M., Shanthikumar, J.G.: *Stochastic convexity and its applications*. *Adv. Appl. Prob.* **20**, 427–446 (1988)
- [10] Skowroński, A.: On some properties of J-convex stochastic processes. *Aequat. Math.* **44**, 249–258 (1992)
- [11] Smolarz, J.: On some functional inequalities connected with quasiconvex functions. *C. R. Math. Rep. Acad. Sci. Canada* **16/6**, 241–246 (1994)

Kazimierz Nikodem
Department of Mathematics and Computer Science
University of Bielsko-Biala
Willowa 2, 43-309 Bielsko-Biala
Poland
e-mail: knikodem@ath.bielsko.pl

Dawid Kotrys
e-mail: dkotrys@gmail.com

Received: February 6, 2014